

All posttests must be submitted online to AAPA.

The CME scoring service is provided at no charge to current Fellow members of AAPA. Others must submit a \$20 scoring fee for each posttest using AAPA's secure server. Payment may be made using Visa, MasterCard, or American Express. To submit this posttest and earn 1 hour of AAPA Category I CME credit, go to www.aapa.org/cme/post-test.html.

TREATING VASOMOTOR SYMPTOMS, PAGE 26

1. The dose of oral conjugated equine estrogens used to alleviate menopausal symptoms is
 - a. 0.625 mg
 - b. 1.0 mg
 - c. 1.25 mg
 - d. 50 mcg
2. The supplement most commonly taken for menopausal symptoms is
 - a. Wild yam
 - b. Vitamin E
 - c. Soy isoflavones
 - d. Ginseng
3. Which of the following side effects has been associated with black cohosh?
 - a. Somnolence
 - b. Headache
 - c. Unpleasant taste
 - d. Photosensitivity
4. Dong quai can potentiate the activity of
 - a. Theophylline
 - b. Digoxin
 - c. Lithium
 - d. Warfarin
5. Women taking evening primrose oil should avoid taking
 - a. Antiepileptic drugs
 - b. Cholesterol absorption inhibitors
 - c. Narcotic analgesics
 - d. Antihypertensives
6. According to studies, hot flash frequency can be reduced by more than 50% with
 - a. Sumatriptan
 - b. Clonazepam
 - c. Clonidine
 - d. Ropinirole

ASEPTIC MENINGITIS, PAGE 54

7. Most cases of aseptic meningitis are caused by
 - a. Influenza
 - b. An enterovirus
 - c. Varicella zoster
 - d. Epstein-Barr virus
8. Which class of medications can replicate the symptoms of aseptic meningitis?
 - a. Antidepressants
 - b. Protease inhibitors
 - c. Antihistamines
 - d. NSAIDs
9. Symptoms similar to those for aseptic meningitis can be caused by immunization for
 - a. Mumps
 - b. *Haemophilus influenzae*
 - c. Pertussis
 - d. Pneumococcal disease
10. On physical examination, aseptic meningitis may be associated with
 - a. Erythematous vesicular oral lesions
 - b. Vesicular eruption on the palms, soles, and buttocks
 - c. Diffuse erythematous, maculopapular rash
 - d. Presence of petechiae or purpuric rash
11. Which finding on analysis of CSF is most indicative of aseptic meningitis?
 - a. $> 1,000$ WBC/mm³
 - b. Protein < 100 mg/dL
 - c. Glucose < 40 mg/dL
 - d. < 0.4 CSF:blood glucose ratio
12. Demyelination syndrome in the setting of aseptic meningitis may result from too rapid administration of
 - a. Aspirin
 - b. Anticonvulsants
 - c. Corticosteroids
 - d. Intravenous fluids

Successful completion of the self-assessment is required to earn Category I CME credit. Successful completion is defined as a cumulative score of at least 70% correct.